

New Zealand Society of Animal Production online archive

This paper is from the New Zealand Society for Animal Production online archive. NZSAP holds a regular annual conference in June or July each year for the presentation of technical and applied topics in animal production. NZSAP plays an important role as a forum fostering research in all areas of animal production including production systems, nutrition, meat science, animal welfare, wool science, animal breeding and genetics.

An invitation is extended to all those involved in the field of animal production to apply for membership of the New Zealand Society of Animal Production at our website www.nzsap.org.nz

[View All Proceedings](#)

[Next Conference](#)

[Join NZSAP](#)

The New Zealand Society of Animal Production in publishing the conference proceedings is engaged in disseminating information, not rendering professional advice or services. The views expressed herein do not necessarily represent the views of the New Zealand Society of Animal Production and the New Zealand Society of Animal Production expressly disclaims any form of liability with respect to anything done or omitted to be done in reliance upon the contents of these proceedings.

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](http://creativecommons.org/licenses/by-nc-nd/4.0/).



You are free to:

Share— copy and redistribute the material in any medium or format

Under the following terms:

Attribution — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

NonCommercial — You may not use the material for [commercial purposes](#).

NoDerivatives — If you [remix, transform, or build upon](#) the material, you may not distribute the modified material.

<http://creativecommons.org.nz/licences/licences-explained/>

Lamb castration and tailing practices in New Zealand: results of a questionnaire

RA Corner-Thomas¹, MB Ferguson², PR Kenyon¹, KJ Stafford¹ and WE Pomroy¹

¹ *International Sheep Research Centre, Institute of Veterinary, Animal and Biomedical Sciences, Massey University, Palmerston North, New Zealand.* ² *The New Zealand Merino Company, Christchurch, New Zealand.*

* *Corresponding author.* Email: r.corner@massey.ac.nz

Abstract

There is a large body of literature that has examined the impacts of castration and tailing on lamb behaviour, stress response, pain and productivity (Fisher et al. 2010; Kent et al. 1993; Mellor et al. 2002; Mellor & Stafford 2000). Recently, Kerslake et al. (2015) investigated lamb tail docking, residual tail length and the reasons behind farmers' choice to dock. There is little information available, however, regarding castration methods and reasons for the choice of leaving lambs entire, castrated or short-scrotumed. A printed questionnaire was distributed to ~12,000 farmers on the Beef + Lamb New Zealand database. The questionnaire contained 30 questions on aspects of management of flystrike, louse control, tailing and castration. Information was sought regarding percentage of lambs tail docked, method used to remove tails, reason for the tailing decision, proportions of male lambs left entire or castrated or short scrotumed, the method used to castrate or short-scrotum, and the reason for choice of male lamb type. A total of 1253 (10%) of questionnaires were returned. Of the farmers that responded, the most commonly used tailing method was a hot iron (61%), followed by rubber rings (31.4%). Lamb growth was the primary reason that farmers gave for leaving lambs entire, and to a lesser degree short-scrotum. The reason behind the decision to castrate was most commonly related to lamb behaviour, however, a number of reasons had similar frequencies (lambs stay cleaner, infertility and meat quality).

Keywords: lamb; castration; tailing docking

Introduction

Routinely in New Zealand, lambs are tail docked and ram lambs castrated or short scrotumed (Stafford 2013). These husbandry procedures are undertaken in order to reduce the incidence of flystrike (tail docking) or to restrict reproduction and facilitate management (castration or short scrotum) (Sutherland 2011). Both castration and tail docking are painful husbandry procedures that are commonly conducted without analgesia and, therefore, are a welfare issue (Kent et al. 1993; Mellor et al. 2002; Mellor & Stafford 2000). These painful procedures and the welfare implications are becoming a focus of attention for consumers, both in New Zealand and world-wide (Phillips et al. 2009). Indeed, it has been suggested that the use of these practices may become a barrier to overseas markets (Hughes 1995).

Recently tail docking, and in particular the length of tail, has been investigated in depth by Kerslake et al. (2015). In addition to collecting data on tail docking practices they examined the reasons for the farmers' choice to tail dock lambs and length of tail remaining (Martin-Collado et al. 2015). To date, however, castration practices used by New Zealand farmers have received little attention. This research, therefore, set out to examine the tailing and castration practices of New Zealand farmers. In addition, the reasons for these practices were also determined.

Materials and methods

In October 2016, a questionnaire was distributed inside the Heartland Sheep issue of Country-wide magazine to ~12,000 farmers on the Beef + Lamb New Zealand database. The questionnaire contained 30 questions on aspects of management of flystrike, louse control, tailing and castration. The first seven questions of the

questionnaire sought information on farmer demographics, i.e., location (regional council area), farmer age, stock units farmed, farm size, numbers of mixed age ewes, mixed-age ewe lambing percentage and sheep breeds. The final six questions related to tailing and castration. The questions were 1) the percentage of lambs that were tailed in 2015, 2) the method used to remove tails, 3) the reason for removing tails or leaving them intact, 4) the proportion of male lambs that were left entire, castrated or short-scrotum, 5) the method used to castrate or short scrotum lambs and 6) the reason for choosing to have entire or castrated or short-scrotum male lambs. Questions 1 to 5 were open allowing respondents to write a response. Question 6 asked respondents to select one or more reasons from a list that included "Lambs are cleaner", "Behaviour", "Lamb growth rate", "Lambs intended to be sold store", "Infertility" and "Meat quality".

Due to the open-ended nature of the responses to question three, they were categorised as follows: "Animal health" = animal health, hygiene, less dags, less flystrike and lambs are cleaner. "Cost" = less labour, reduced costs. "Management" = ease of management, saves time, easier shearing/crutching, less dagging and easier breeding. "Historical" = always done it and common sense. "Lamb type" = sex of lamb, age of lamb, terminal-sired lambs, time of slaughter. "Buyer/works requirements" = works requirements or buyer requirements. "Aesthetics" = looks, unnecessary, tidy.

A total of 1253 (10%) questionnaires were returned of which 1224 were included in the analyses. Questionnaires were excluded if no response was provided for either percentage of lambs tailed or percentage of lambs that were entire, castrated or short scrotumed.

Statistical analysis

All statistical analyses were conducted using SAS (SAS Institute, Cary, NC, USA). The data were summarised by determining the frequency of respondents that used each tailing or castration method. For each method the frequency of farmers that gave each reason (behaviour, lambs are cleaner, lamb growth rate, infertility, meat quality or lambs sold store) was determined. The summarised data was then analysed using a generalised model with a binomial distribution. For the analysis of reason for tail docking, the outcome variable was the percentage docked (tailed 95% or more vs. tailed less than 95%). The model for castration reason included the fixed effect of male lamb type (entire, castrate or short scrotum).

Results

Demographics

The farmers that returned completed surveys and had usable data (n=1224) represented farms with a total combined effective area of 890,000 ha and 5.8 million stock units with a mean flock size of 2,280 ewes, an effective area of 751 ha and a lambing percentage (lambs tailed per ewe put to the ram) of 159%. The median age of the respondents was “50 to 59 years” of age with less than 10% of respondents being under 40 years of age. The majority of farms were located in the North Island (n=586), followed by the south of the South Island (n=329) and the north of the South Island (n=241) with the remainder (n=68) not providing location information. The vast majority of respondents (n=1107) farmed strong-wool breeds such as Romney, Coopworth and Perendale and crosses containing these breeds. Only a small number had either medium (n=35) or fine (n=38) wool breeds or shedding sheep (n=23). A small number of farmers did not indicate the breeds they farmed (n=21).

Tailing

Of the questionnaires returned 1089 (89%) respondents reported the proportion of lambs that they tail docked. Tail docking was common with 757 (62%) that docked 100% of their lambs and 995 (91%) that docked more than 95%. A small number of respondents (n=27, 2.5%) did not dock any lambs. A total of 1186 respondents, provided information on the tail docking method they used. The majority used only one method to tail dock lambs, using either a cauterising iron (n=721, 61%), rubber ring (372, 31.4%) or knife (n=10, 0.8%). A small number of respondents (n=83, 7.0%) used two methods, the most common being the cauterising iron and rubber ring (n=43, 3.6%) and rubber ring and knife (n=36, 3%). Although only a small number of farmers used a knife to castrate all lambs, 51 (4%) used a knife in conjunction with another method.

The reasons all respondents gave for their decision to tail dock lambs were predominantly related to animal health (58%, 95% CI = 56-60%) and ease of management (30%,

Table 1 The frequency (n) for each reason given for the choice to remove a lamb’s tail or leave it intact and a comparison of the percentage (back-transformed mean with 95% confidence intervals) of respondents that gave each reason between farmers that did, or did not, tail dock 95% or more of their lambs.

	n	Tail docked 95% or more of lambs		No vs. Yes
		No (n=94)	Yes (n=995)	
Health	1400	38.6 (31.6 - 46.1) ^d	59.7 (57.6 - 61.9) ^c	< 0.05
Management	722	15.8 (11.1 - 22.1) ^b	30.8 (28.8 - 32.9) ^d	< 0.05
Lamb type	77	27.5 (21.3 - 34.7) ^c	1.2 (0.8 - 1.8) ^b	< 0.05
Other	63	9.4 (5.8 - 14.7) ^b	1.9 (1.4 - 2.6) ^c	< 0.05
Cost/labour	37	3.5 (1.6 - 7.6) ^a	1.3 (0.9 - 1.9) ^b	0.03
Buyer/works	62	2.9 (1.2 - 6.8) ^a	2.5 (1.9 - 3.3) ^c	ns
Welfare	32	1.8 (0.6 - 5.3) ^a	1.4 (1 - 2) ^b	ns
Historical	15	0.6 (0.1 - 4) ^a	0.5 (0.3 - 1) ^a	ns
Aesthetics	13	---	0.5 (0.3 - 1) ^a	n/a

^{abcde} superscripts that differ within columns indicate that percentages were significantly different (P<0.05).

95% CI = 28-32%). Of the respondents that tail docked 95% of their lambs, a greater percentage gave reasons relating to animal health and ease of management (Table 1). For those that tailed less than 95% a greater (P<0.05) percentage of respondents gave reasons relating to cost/labour and lamb type.

Entire or Castrated or Short scrotum male lambs

Based on the proportions reported by each respondent, the majority of lambs were short scrotum (42%) with similar proportions left entire or castrated (29% and 25%, respectively). Of the 1224 respondents, 1191 (97%) reported the proportions of male lambs that were left entire, castrated or short scrotum. From these proportions it was calculated that 35% (n=423) of respondents castrated some or all lambs, 48% (n=587) short scrotum and 52% (n=638) left lambs entire. The majority either only castrated (n=196, 16.5%), short scrotum (n=310, 26.0%) or left lambs entire (n=269, 22.6%). Some farmers, however, had more than one type of male lamb, for example, 15.9% (n=189) had entire and short-scrotum lambs, 9.8% (n=117) had entire and castrated lambs, 4.0% (n=47) had castrated and short-scrotum lambs. There were a small number of farmers that had all three male lambs types (n=63, 5.3%).

When asked for reasons for choosing to leave lambs entire, lamb growth rate was cited by the majority of farmers (Table 2). Similar percentages (P>0.05) of respondents gave “Lambs sold store”, “Meat quality” and “Lambs are cleaner” as reasons for their choice to leave lambs entire. The decision to castrate was most commonly chosen for lamb behaviour reasons. Similar percentage of respondents choose “Cleaner”, “Infertility” and “Meat quality” as the reason for their decision to castrate. The most commonly cited reason for short scrotum lambs was “Lamb growth rate”. Similar percentages (P>0.05) of farmers cited “Behaviour”, “Lambs are cleaner” and “Infertility” as reasons for their choice.

Table 2 The frequency (n) for each reason for the choice to have entire, castrated and short scrotum lambs and a comparison of each reason (back-transformed mean with 95% confidence intervals) given by farmers for having entire (Ent), castrated (Cast) or short-scrotum lambs (SS).

Reason given	Male lamb type						P-value		
	n	Entire	n	Castrate	n	Short scrotum	Cast vs. SS	Cast vs. Ent	SS vs. Ent
Behaviour	42	6.1 (4.6 - 8.2) ^b	266	24.1 (21.6 - 26.7) ^d	302	16.2 (14.6 - 18) ^b	**	**	**
Lambs are cleaner	65	9.4 (7.4 - 11.8) ^c	191	17.3 (15.2 - 19.6) ^c	317	17 (15.4 - 18.8) ^b	ns	**	**
Lamb growth rate	423	59.9 (56.2 - 63.6) ^d	98	8.9 (7.3 - 10.7) ^a	466	25.1 (23.1 - 27.1) ^c	**	**	**
Infertility	25	3.5 (2.4 - 5.2) ^a	193	17.5 (15.3 - 19.8) ^c	339	18.2 (16.5 - 20.1) ^b	ns	**	**
Meat quality	71	10.4 (8.3 - 12.9) ^c	216	19.5 (17.3 - 22) ^c	232	12.5 (11.1 - 14.1) ^a	**	**	ns
Lambs sold store	73	10.7 (8.6 - 13.2) ^c	142	12.8 (11 - 14.9) ^b	204	11 (9.6 - 12.5) ^a	ns	ns	ns

^{abcd} within columns superscripts that differ indicate that percentages were significantly different ($P < 0.05$).

ns = not significant ($P > 0.05$), * = $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Discussion

The aim of the current study was to investigate the tailing and castration practices of New Zealand sheep farmers and to determine the reasons behind the use of each method. The response rate in the current study was relatively low at 10%, however, due to the questionnaire being sent to every sheep farmer New Zealand, it generated more than 1,200 responses which has created a significant database of information on farmer practices.

The farmers that returned completed questionnaires farmed an average effective area similar to those that completed previous questionnaires conducted by our group (862 ha for a questionnaire in 2014 and 607 ha in 2012; Corner-Thomas et al. 2015; 2016). The farmers that completed the questionnaire had larger farms and had higher lambing percentages than the average New Zealand farm which has an effective size of 250 ha with an average flock size of 1,588 and a lambing percentage of 127% (Beef + Lamb NZ 2016). The age distribution of the farmers that returned a completed questionnaire was similar to that reported in the questionnaires conducted in 2012 and 2014 with the median age of “50 to 59 years” of age (Corner-Thomas et al. 2016).

Tail docking was a common practice amongst the farmers that completed the questionnaire with the majority (62%) docking 100% of lambs with a small number (2.5%) that did not tail dock any lambs. These results contrast with those of Kerslake et al. (2015) who reported that 6% of respondents tail docked 100% of their lambs and 1.4% did not tail dock any. It is possible that the differences in the findings of these studies were the result of the differences in the method of distribution of the questionnaires (postal vs. online) resulting in a different demographic of farmers (Casey et al. 2016; Corner-Thomas et al. 2017). The tail docking methods used by the farmers in the current study, however, were similar to those reported by Kerslake et al. (2015) with approximately 60% of farmers using a cauterising iron and 30% using a rubber ring. Somewhat surprising was that 4% of respondents tail docked lambs with a knife (surgical amputation). Although not actively discouraged, the animal-welfare painful-husbandry procedures code of animal welfare (Anon. 2005) noted

that surgical techniques are associated with greater risks of bleeding and infection.

The primary reason cited for tail docking lambs was animal health, which included responses relating to minimising flystrike, minimising dags, having cleaner lambs and animal health. Kerslake et al. (2015) and Fisher et al. (2006) also reported a similar result with farmers that docked lambs ranking minimising flystrike followed by reducing dags and reducing crutching costs as the three most important reasons for their docking practices. It is perhaps unsurprising that minimising flystrike and dags are often given as a reason to tail dock given that docking has been shown to reduce the incidence in flystrike (French et al. 1994; Webb-Ware et al. 2000).

In the current study, 35% of farmers had more than one male lamb type (entire, short-scrotum and castrate) on their farm. This finding is in agreement with that of Tarbotton et al. (2002) who reported 30% of farmers had more than one type of male lamb. The variation in the percentages of each male lamb type are likely due to the intended final use of the animal. Lambs that will be sold to slaughter soon after weaning are more likely to be left entire, as management of fertility is not required, whereas lambs kept on farm into the breeding season requires management of fertility and behaviour (Johnson et al. 2017). Many farmers (65%), however, chose to have only one male lamb type.

The reasons cited for the choice of entire, short scrotum or castrated lambs differed between farmers. Farmers that left some or all lambs entire or short scrotum commonly gave lamb growth rate as a reason for their choice, whereas, farmers that castrated lambs commonly cited lamb behaviour. This finding is similar to that of Tarbotton et al (2002) who reported that the most common reason given to cryptorchid or leave lambs entire was better growth, however, for castration the most commonly given reason was convenience (fast, easier). In the questionnaire conducted by Tarbotton et al. (2002), behaviour was only given by 8% of respondents as a reason to castrate.

Conclusion

Tail docking and castration practices of New Zealand farmers reflect the diverse nature of the sheep-production

systems, with farmers utilising a range of tailing and castration practices. The reasons for the choice of tailing practice primarily centred on animal health and, in particular, reducing dags and risk of flystrike. Lamb growth was the most commonly cited reason for leaving lambs entire and short scrotummed, however, behavioural reasons were most common for castration.

Acknowledgements

This research was funded by The New Zealand Merino Company.

References

- Anon. 2005. Animal welfare (painful husbandry procedures) code of welfare. In: Committee NAWA ed. Code of Welfare No. 7 Wellington, New Zealand, Ministry for Primary Industries, Government of New Zealand. Pp. 36.
- Beef + Lamb NZ 2016. Compendium of New Zealand farm facts. Wellington, New Zealand, Beef + Lamb New Zealand. 28 p.
- Casey M, Meikle A, Kerr G, Stevens D 2016. Social media—a disruptive opportunity for science and extension in agriculture? In: Thom E ed. Hill Country Symposium. Pp. 53-60.
- Corner-Thomas R, Kenyon P, Morris S, Ridler A, Hickson R, Greer A, Logan C, Blair H 2017. Farmer perceptions of the relative usefulness of information providers and technology transfer methods. *New Zealand Journal of Agricultural Research* 60: 245-262.
- Corner-Thomas RA, Kenyon PR, Morris ST, Ridler AL, Hickson RE, Greer AW, Logan CM, Blair HT 2015. Influence of demographic factors on the use of farm management tools by New Zealand farmers. *New Zealand Journal of Agricultural Research* 58: 412-422.
- Corner-Thomas RA, Kenyon PR, Morris ST, Ridler AL, Hickson RE, Greer AW, Logan CM, Blair HT 2016. Brief communication: The use of farm-management tools by New Zealand sheep farmers: Changes with time. *Proceedings of the New Zealand Society of Animal Production* 76: 78-80.
- Fisher M, Agnew R, Thomas A, Scobie D 2006. Tail docking sheep—why is it undertaken, what are the issues, and what costs and benefits are associated with it and the alternative of breeding short-tailed animals. Ministry of Agriculture and Forestry Operational Research, Final report Contract AWE212-Docking sheep.
- Fisher MW, Bray AR, Johnstone PD 2010. Implications of removing or altering the testicles of ram lambs on the financial returns from carcasses. *New Zealand Journal of Agricultural Research* 53: 135-143.
- French NP, Wall R, Morgan KL 1994. Lamb tail docking: A controlled field study of the effects of tail amputation on health and productivity. *Veterinary Record* 134: 463-467.
- Hughes D 1995. Animal welfare: The consumer and the food industry. *British Food Journal* 97: 3-7.
- Johnson PL, Stuart AD, Mackie A 2017. Brief communication: High ultimate pH in short-scrotum male lambs processed during the breeding season. *Proceedings of the New Zealand Society of Animal Production*. Rotorua. Pp. 194-196.
- Kent J, Molony V, Robertson I 1993. Changes in plasma cortisol concentration in lambs of three ages after three methods of castration and tail docking. *Research in Veterinary Science* 55: 246-251.
- Kerslake JI, Byrne TJ, Behrent MJ, Maclennan G, Martin-Collado D 2015. The reasons farmers choose to dock lamb tails to certain lengths, or leave them intact. *Proceedings of the New Zealand Society of Animal Production* 75: 210-214.
- Martin-Collado D, Byrne TJ, Amer PR, Behrent MJ, Maclennan G, Kerslake JI 2015. Analysing hidden patterns of farmers' preferences for farm performance characteristics that may be related to tail-docking practice decisions. *Proceedings of the New Zealand Society of Animal Production* 75: 205-209.
- Mellor D, Stafford K, Todd S, Lowe T, Gregory N, Bruce R, Ward R 2002. A comparison of catecholamine and cortisol responses of young lambs and calves to painful husbandry procedures. *Australian Veterinary Journal* 80: 228-233.
- Mellor DJ, Stafford KJ 2000. Acute castration and/or tailing distress and its alleviation in lambs. *New Zealand Veterinary Journal* 48: 33-43.
- Phillips CJC, Wojciechowska J, Meng J, Cross N 2009. Perceptions of the importance of different welfare issues in livestock production *Animal* 3: 1152-1166.
- Stafford KJ 2013. *Animal welfare in New Zealand*. Cambridge, New Zealand, New Zealand Society of Animal Production 204 p.
- Sutherland MA 2011. Painful husbandry procedures and methods of alleviation: A review. *Proceedings of the New Zealand Society of Animal Production* 71: 189-194.
- Tarbotton IS, Bray AR, Wilson JA 2002. Incidence and perceptions of cryptorchid lambs in 2000. *Proceedings of the New Zealand Society of Animal Production* 62: 334-336.
- Webb-Ware JK, Vizard AL, Lean GR 2000. Effects of tail amputation and treatment with an albendazole controlled-release capsule on the health and productivity of prime lambs. *Australian Veterinary Journal* 78: 838-842.