

New Zealand Society of Animal Production online archive

This paper is from the New Zealand Society for Animal Production online archive. NZSAP holds a regular annual conference in June or July each year for the presentation of technical and applied topics in animal production. NZSAP plays an important role as a forum fostering research in all areas of animal production including production systems, nutrition, meat science, animal welfare, wool science, animal breeding and genetics.

An invitation is extended to all those involved in the field of animal production to apply for membership of the New Zealand Society of Animal Production at our website www.nzsap.org.nz

[View All Proceedings](#)

[Next Conference](#)

[Join NZSAP](#)

The New Zealand Society of Animal Production in publishing the conference proceedings is engaged in disseminating information, not rendering professional advice or services. The views expressed herein do not necessarily represent the views of the New Zealand Society of Animal Production and the New Zealand Society of Animal Production expressly disclaims any form of liability with respect to anything done or omitted to be done in reliance upon the contents of these proceedings.

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](http://creativecommons.org/licenses/by-nc-nd/4.0/).



You are free to:

Share— copy and redistribute the material in any medium or format

Under the following terms:

Attribution — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

NonCommercial — You may not use the material for [commercial purposes](#).

NoDerivatives — If you [remix, transform, or build upon](#) the material, you may not distribute the modified material.

<http://creativecommons.org.nz/licences/licences-explained/>

BRIEF COMMUNICATION: Variability in growth rates of goat kids on 16 New Zealand dairy goat farms

LE Deeming^{1,2}, NJ Beausoleil², KJ Stafford², JR Webster¹, and G Zobel^{1*}

¹AgResearch Ltd., Ruakura Research Centre, 10 Bisley Road, Private Bag 3123, Hamilton 3214, NZ. ²Institute of Veterinary, Animal and Biomedical Sciences, Massey University, Private Bag 11 222, Palmerston North 4442, NZ

*Corresponding author. Email: gosia.zobel@agresearch.co.nz

Keywords: dairy goats; weaning; average daily gain; management

Introduction

Average daily weight gains (ADG) in early life are associated with health, welfare and future production potential of dairy animals. While this has yet to be established for goats, research in cows demonstrates a positive relationship between growth in early life and milk production (Shamay et al. 2005, Soberon et al. 2012). Early growth rates are largely determined by the management practices from birth onwards. It is therefore unsurprising that large differences in ADG under different management systems have been reported. In lambs, Mahgoub et al. (2000) reported a large range in ADG (84-154 g/day). Similarly, in calves, Bartlett et al. (2006) reported ADGs ranging from 251 g/day to 703 g/day. The ADGs reported in goats have not been as variable (152-170 g/day: Galina et al. 1995; 167-173 g/day: Goetsch et al. 2001); however, these were small, controlled studies. To our knowledge, no research has yet quantified variability in the ADG of goat kids on a large, multi-farm scale. Therefore, the aim of this study was to describe the variation in growth rates on multiple dairy goat farms in the Waikato region of NZ, providing a benchmark for dairy goat farmers.

Materials and methods

This study was approved by AgResearch Ltd, Ruakura Animal Ethics Committee (13478). Sixteen Dairy Goat Cooperative (DGC) farmers from the Waikato region (NZ) agreed to participate in an on-going three year study. Between May and August 2015, 1269 Saanen Alpine cross kids (between 70 and 83 kids/farm) were enrolled. Kids were weighed four times: at enrolment (between 24 and 48 h after birth), two weeks after enrolment, at weaning, and two weeks after weaning. Kids were weighed near mid-day to minimise the effect of gut fill differences. For each kid, ADG was calculated between each weigh point. Descriptive data were summarised by farm (PROC UNIVARIATE, SAS 9.2), and are presented as means \pm SD (n = 16). Where appropriate, individual kid data is presented and discussed (n = 1269).

Results and discussion

Table 1 summarises the mean weights at each time point on the 16 farms. The weights were variable across farms, even at the early weigh points. For example, the smallest kid at enrolment was 1.4 kg and the largest was

6.3 kg. When kids were weighed two weeks later, the lowest weight recorded was 2.9 kg and the highest was 13.5 kg. While birth weight will account for some weight gain variation early in life (calves: Donovan et al. 1998), colostrum management is likely a greater contributor (goat kids: Abdou et al. 2014; lambs: Massimini et al. 2006). How colostrum was delivered (e.g. dam, bottle), timing of delivery (e.g. immediately vs. up to 24 h after birth) and type of colostrum (e.g. caprine, bovine, bovine-based replacement product) differed on the 16 farms. Adequate colostrum intake after birth is known to be important in developing a young ruminant's immune system (Hurley & Theil 2011); without this development, disease risk increases (O'Brien & Sherman 1993), which subsequently may decrease growth rates (Massimini et al. 2006).

Weaning weights were variable, even among farms with similar weaning ages. For example, the farm with the lowest weaning weights (14.7 \pm 2.2 kg) weaned at 10.0 \pm 0.9 weeks old, while the farm with the highest weaning weights (24.2 kg \pm 3.0 kg) weaned at 11.2 \pm 0.3 weeks old. This suggests that factors other than age at weaning impact growth at this critical time. Overall energy intake pre-weaning (Bartlett et al. 2006), dictated by the quality and quantity of feed provided, is a determinant of young ruminant growth (Mahgoub et al. 2000). Most farms offered milk *ad libitum*, while two restricted feedings to twice a day; types of milk fed included whey, goat milk, and cow milk powders, as well as fresh goat and cow milk. All farmers provided hard feeds, but the type provided, as well as the timing of its provision varied amongst farms.

From enrolment to weaning, ADG (Table 1) were consistent with pre-weaning ADG previously reported (e.g. 167-173 g/day: Goetsch et al. 2001). Interestingly, the variability across farms was more pronounced in the period between weaning and two weeks after weaning (Table 1). Five farms achieved consistent ADG after weaning compared to before weaning, and three of these farms achieved post-weaning ADGs > 200 g/day (Figure 1). However, ADG on the remaining 11 farms decreased after weaning. Of these, five farms experienced large decreases in ADG, achieving growth of < 100 g/day, and one farm had a negative ADG. While this latter farm reported cases of pneumonia, reduced ADG or even weight loss occurring after weaning is well documented regardless of health status (calves: Sweeney et al. 2010; goat kids: Ugur et al. 2004).

Figure 1 Mean (\pm SD) average daily gain (ADG) of goat kids ($n = 80 \pm 3$ per farm) on 16 New Zealand dairy goat farms from enrolment until weaning (light grey) and weaning until 2 weeks after weaning (dark grey).




Table 1 Mean (\pm SD) age, weight and average daily gain (ADG) of goat kids on New Zealand dairy goat farms ($n=16$) at four weigh points.

	Weigh point			
	Enrolment	Two weeks after enrolment	Weaning	Two weeks after weaning
Age (days)	1.6 \pm 0.4	15.9 \pm 4.0	87.2 \pm 13.0	101.3 \pm 13.0
Weight (kg)	3.7 \pm 0.6	6.2 \pm 0.9	19.2 \pm 2.7	21.3 \pm 3.2
ADG (g/day) ¹	-	177 \pm 32	180 \pm 40	137 \pm 66
ADG (g/day) ²	-	-	184 \pm 24	177 \pm 23

¹calculated between each weigh point

²calculated relative to enrolment weigh point

This growth lag may be a consequence of stress induced by coping with a new diet (Magistrelli et al. 2013), or due to inadequate rumen development from insufficient hard feed consumption prior to weaning (Khan et al. 2007).

The findings of this study indicate that there is large variability in growth rates of dairy goat kids on NZ farms. Further analysis will aim to quantify how farm-specific factors (e.g. weight at birth, colostrum/milk feeding management, weaning management), may be causing these differences in ADG. It should be noted that one farm consistently achieved ADG > 200 g/day from birth through to two weeks after weaning. This provides a benchmark for NZ dairy goat farmers and suggests that high, stable weight gains, immediately following weaning, are possible.

Acknowledgements

Funding for the research and a student stipend is provided by the NZ Ministry of Business, Innovation & Employment (MBIE), and the Dairy Goat Cooperative (NZ) Ltd (DGC). Special thanks goes to the 16 participating DGC farmers.

References

Abdou H, Marichatou H, Beckers JF, Dufrasne I, Issa M, Hornick JL 2014. Effect of bovine colostrum intake on growth, reproductive parameters and survival in red kids. *Journal of Animal Physiology and Animal Nutrition* 98: 845-852.

Bartlett K, Mckeith F, Vandehaar M, Dahl G, Drackley J 2006. Growth and body composition of dairy calves fed milk replacers containing different amounts of protein at two feeding rates. *Journal of Animal Science* 84: 1454-1467.

Donovan GA, Dohoo IR, Montgomery DM 1998. Calf and disease factors affecting growth in female Holstein calves in Florida, USA. *Preventative Veterinary Medicine* 33: 1-10.

Galina MA, Palma JM, Pacheco D, Morales R 1995. Effect of goat milk, cow milk, cow milk replacer and partial substitution of the replacer mixture with whey on artificial feeding of female kids. *Small Ruminant Research* 17: 153-158.

Goetsch AL, Detweiler G, Sahlu T, Dawson LJ 2001. Effects of different management practices on preweaning and early postweaning growth of Alpine kids. *Small Ruminant Research* 41: 109-116.

Hurley WL, Theil PK 2011. Perspectives on immunoglobulins in colostrum and milk. *Nutrients* 3: 442-474.

Khan MA, Lee HJ, Lee WS, Kim HS, Ki KS, Hur TY, Suh GH, Kang SJ, Choi YJ 2007. Structural growth, rumen development, and metabolic and immune responses of holstein male calves fed milk through step-down and conventional methods. *Journal of Dairy Science* 90: 3376-3387.

Magistrelli D, Aufy AA, Pinotti L, Rosi F 2013. Analysis of weaning-induced stress in Saanen goat kids. *Journal of Animal Physiology and Animal Nutrition* 97: 732-739.

Mahgoub O, Lu CD, Early RJ 2000. Effects of dietary energy density on feed intake, body weight gain and carcass chemical composition of Omani growing lambs. *Small Ruminant Research* 37: 35-42.

Massimini G, Britti D, Peli A, Cinotti S 2006. Effect of passive transfer status on preweaning growth performance in dairy lambs. *Journal of the American Veterinary Medical Association* 229: 111-115.

O' Brien JP, Sherman DM 1993. Serum immunoglobulin concentrations of newborn goat kids and subsequent kid survival through weaning. *Small Ruminant Research* 11: 71-77.

Shamay A, Werner D, Moallem U, Barash H, Bruckental I 2005. Effect of Nursing Management and Skeletal Size at Weaning on Puberty, Skeletal Growth Rate, and Milk Production During First Lactation of Dairy Heifers. *Journal of Dairy Science* 88: 1460-1469.

Soberon F, Raffrenato E, Everett RW, Van Amburgh ME 2012. Preweaning milk replacer intake and effects on long-term productivity of dairy calves. *Journal of Dairy Science* 95: 783-793.

Sweeney BC, Rushen J, Weary DM, De Passillé AM 2010. Duration of weaning, starter intake, and weight gain of dairy calves fed large amounts of milk. *Journal of Dairy Science* 93: 148-152.

Ugur F, Savas T, Dosay M, Karabayır A, Atasoglu C 2004. Growth and behavioral traits of Turkish Saanen kids weaned at 45 and 60 days. *Small Ruminant Research* 52: 179-184.