

## New Zealand Society of Animal Production online archive

This paper is from the New Zealand Society for Animal Production online archive. NZSAP holds a regular annual conference in June or July each year for the presentation of technical and applied topics in animal production. NZSAP plays an important role as a forum fostering research in all areas of animal production including production systems, nutrition, meat science, animal welfare, wool science, animal breeding and genetics.

An invitation is extended to all those involved in the field of animal production to apply for membership of the New Zealand Society of Animal Production at our website [www.nzsap.org.nz](http://www.nzsap.org.nz)

[View All Proceedings](#)

[Next Conference](#)

[Join NZSAP](#)

The New Zealand Society of Animal Production in publishing the conference proceedings is engaged in disseminating information, not rendering professional advice or services. The views expressed herein do not necessarily represent the views of the New Zealand Society of Animal Production and the New Zealand Society of Animal Production expressly disclaims any form of liability with respect to anything done or omitted to be done in reliance upon the contents of these proceedings.

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](http://creativecommons.org/licenses/by-nc-nd/4.0/).



You are free to:

**Share**— copy and redistribute the material in any medium or format

Under the following terms:

**Attribution** — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

**NonCommercial** — You may not use the material for [commercial purposes](#).

**NoDerivatives** — If you [remix, transform, or build upon](#) the material, you may not distribute the modified material.

<http://creativecommons.org.nz/licences/licences-explained/>

## BRIEF COMMUNICATION: High ultimate pH in short-scrotum male lambs processed during the breeding season

PL Johnson<sup>a\*</sup>, AD Stuart<sup>b</sup> and A Mackie<sup>b</sup>

<sup>a</sup>AgResearch Invermay, Puddle Alley, Mosgiel, New Zealand; <sup>b</sup>Alliance Group Limited, PO Box 845, Invercargill

\*Corresponding author. Email: tricia.johnson@agresearch.co.nz

**Keywords:** lamb; sex; meat quality; pH

### Introduction

Historically in New Zealand male lambs were castrated at the time of docking, however, there has been a move to instead create short-scrotum males, in which the testes are pushed within the body cavity and the scrotum removed (Johnson et al. 2007). This results in short-scrotum males, having similar physiological characteristics to entire ram lambs, except that they are almost always infertile and, therefore, do not present a management risk if they are accidentally mixed with ewes or ewe lambs during the mating season. The normal ultimate pH of lamb meat is considered to be 5.50 to 5.80, with values above 5.8 and in particular above 6.0 considered high, and resulting in negative meat quality outcomes. There have been a number of studies that have investigated whether or not sex differences in the ultimate pH of lamb exist. The results are not always consistent with some studies reporting elevated ultimate pH in non-castrated male lambs (Johnson et al. 2005; Bain et al. 2009), whilst others have shown no differences (Kerslake et al. 2012; Schreurs 2013). In commercial farming operations there is a mix as to how farmers manage their lambs of different sexes during the finishing period prior to slaughter, and in many farming operations this period coincides with the on-set of the breeding season.

Given the most recent papers in this area have concluded that there is no sex-effect on ultimate pH, the purpose of this paper is to report on data collected from a trial that set out to invoke high ultimate pH (to investigate potential live-animal predictors of high ultimate pH) in short-scrotum male lambs during the breeding season.

### Materials and methods

#### Animals Used

Animals for this study were sourced from a commercial property in Southland, New Zealand in mid-April 2015. Ethical approval for the trial was obtained from the AgResearch Invermay Animal Ethics Committee (Application: AGR13507).

The experimental design required to investigate live-animal predictors for *M. longissimus lumborum* ultimate pH required a sample size of 22 animals with high ultimate pH. In previously obtained data sets, it was determined that the proportion of animals with ultimate pH above 5.8 could be as low as 10% (Johnson et al. 2015), and therefore a data set consisting of 220 animals was generated.

A group of short-scrotum male lambs of maternal-composite breeding with a minimum live weight of 40kg were placed in a paddock adjacent to ewes and entire rams during the autumn mating period with the assumption that at least a proportion of the ewes would be cycling on any given day. The lambs were grazed next to the ewes from April 8th for 8 days then held overnight in the yards prior to trucking to the abattoir. The lambs were loaded onto a truck at approximately 9.30am and transported for approximately 60 minutes to the Alliance Lorneville plant (near Invercargill). The lambs were held overnight at the abattoir and slaughtered on the morning of April 17th.

#### Measurements made

From their time of arrival at the abattoir, the lambs and subsequent carcasses were processed according to commercial procedures of Alliance Group Ltd. And underwent accelerated conditioning and ageing processes (Chrystall et al. 1989) designed to maximise meat-quality outcomes. The ultimate pH was measured 72 hours after slaughter by insertion of a pH meat probe, attached to a WTW pH330i pH Meter calibrated for temperature, into the *M. longissimus lumborum* muscle.

Additional carcass data as routinely generated by Alliance Ltd. included carcass weight, and measurements reported from their VIAscan two-dimensional imaging system (Hopkins et al. 2004) which included VIAscan GR (an estimate of carcass fatness), and VIAscan estimates of the lean meat yield in the leg, loin and shoulder regions, summed to give total lean meat yield.

#### Analysis

Summary statistics were prepared, and the distribution of the ultimate pH values plotted using the box plot procedure in the statistical programme R. The relationship between ultimate pH and the carcass data set was assessed using the correlation procedure in SAS (SAS, 2011).


### Results and discussion

In studies where the sex contrast has been between ewe and non-castrated male lambs, significant differences in ultimate pH have been reported (Johnson et al. 2005; Bain et al. 2009), whilst in studies that have compared non-castrated lambs to castrated lambs no differences have been observed (Kerslake et al. 2012; Schreurs 2013). In commercial farming operations, grazing and finishing of lambs occurs with lambs in mixed- or single-sex groups.

**Table 1** Ultimate pH and carcass characteristics of 225 maternal-composite short-scrotum lambs slaughtered in mid-April

	n	Average ± StdDev	Range (Min- Max)
Ultimate <i>M. longissimus lumborum</i> pH measured three days post mortem	224	5.98 ± 0.31	5.50-7.24
Carcass weight (kg)	225	18.7 ± 1.49	14.9-22.8
VIAscan GR (mm; measure of fatness)	219	4.0 ± 2.07	1-10
VIAscan lean meat yield leg (%)	219	22.5 ± 1.2	19.7-25.8
VIAscan lean meat yield loin (%)	219	15.3 ± 0.8	13.3-17.5
VIAscan lean meat yield shoulder (%)	219	17.1 ± 0.8	14.8-19.4
VIAscan lean meat yield total (%)	219	55.0 ± 2.3	50.0- 60.7

**Figure 1** Distribution of pH values from 224 non-castrated male lambs slaughtered in mid-Autumn. The box represents 25-75% of the values, with the solid middle bar the median. The top whisker represents two times the interquartile range, with the circles representing outliers. The bottom whisker represents the bottom 25% of values.



In this study, the impact of non-castrated males running adjacent to mature ewes that were running with rams during their breeding season is reported. During mustering, the night before trucking, the shepherd made a notable observation that although the lambs were in a paddock adjacent to the yards, it took him over 40 minutes to muster them into the yards as a proportion of the flock continually doubled back to the fence line adjacent to the ewes. Additionally, mounting behaviour continued in the pre-trucking period.

The summary data for the carcasses is provided in Table. 1. From this summary it can be observed that given the normally acceptable lamb *M. longissimus lumborum* ultimate pH range is 5.5 to 5.8, and the range in the current study was 5.50 to 7.24 (Figure 1) with an average value of 5.98 indicates that the overall ultimate pH of the mob was very high. Over 45% of the lambs had ultimate pH of 6.0 or greater.

Minimum levels of glycogen are required to achieve a full post-slaughter drop in muscle pH (Immonen &

Puolanne 2000). A low glycogen concentration with the post-mortem muscle will reduce production of lactic acid reducing the drop in pH in the muscle, such that it does not achieve the expected range of 5.5 to 5.8. Once pubertal, ram lambs together with adult rams are responsive to the behavioural and physiological characteristics of oestrus ewes, and studies have shown that adult rams in mating mob situations can lose up to 12% of their body weight as a result of decreased feeding and increased activity, activities that were observed in the lambs in this study. The impact of reduced feeding, increased activity and sexual activity on reducing muscle glycogen levels pre-slaughter have been reviewed by Carragher and Matthews (1996), and are considered to be cumulative.

There was no correlation between pH and the carcass traits ( $P > 0.05$ ) for all trait combinations, with no Pearson correlation co-efficient greater than 0.08. This indicates that no attribute of the animal in terms of its size or level of maturity as indicated by carcass fatness were predictive of whether or not the animal would exhibit high pH.

This data set has demonstrated that high ultimate pH can be an issue in non-castrated male lambs, retained through the mating season. As such, management practices need to ensure that non-castrated male lambs are well-segregated from mature and pubertal ewes in order to prevent muscle glycogen depletion and elevated ultimate pH. This should also include ensuring that no cull ewe lambs are accidentally incorporated into non-castrated male finishing mobs.

**Acknowledgements**

Funding for the parent trial was sourced from the AgResearch Curiosity Fund. The authors would like to acknowledge the staff at the Alliance Lorneville plant for their help in data collection.

**References**

Bain WE, McLean NJ, Young EA, Johnson PL 2009. Effect of sex on colour and pH of chilled lamb loins. Proceedings of the New Zealand Society of Animal Production 69: 101–103.

Carragher JF, Matthews LR 1996. Animal behaviour and stress: impacts on meat quality. Proceedings of the New Zealand Society of Animal Production 56: 162-166.

Chrystall, BB, Devine CE, Longdill GR, Gill CO, Swan JE, Petersen GV 1989. Trends and developments in meat processing. Pages 185–207 In Purchas RW, Butler-Hogg BW and Davies AS ed. Meat production and processing, Occasional Publication No. 11. N.Z. Soc. Anim. Prod. Inc., Hamilton, New Zealand.

Hopkins DL, Safari E, Thompson JM, Smith CR 2004. Video image analysis in the Australian meat industry - precision and accuracy of predicting lean meat yield in lamb carcasses. Meat Science 67: 269-274.

- Immonen, K, Puolanne E 2000. Variation of residual glycogen-glucose concentration at ultimate pH values below 5.75. *Meat Science* 55: 279-283.
- Johnson, PL, Purchas, RW, McEwan, JC, Blair, HT 2005. Carcass composition and meat quality differences between pasture-reared ewe and ram lambs. *Meat science* 71: 383-391.
- Johnson, PL, Kenyon, PR, Burnham, DL, West, DM 2007. To wether – short scrotum or leave lambs as entire rams? Revisiting an old question using new genetics. *Proceedings of the New Zealand Society of Animal Production* 67: 44-47.
- Johnson PL, Charteris, A, Tate, ML, Dodds, KG 2015. BRIEF COMMUNICATION: Genetic parameters for meat traits assessed in the Headwaters New Zealand Progeny Test, *Proceedings of the New Zealand Society of Animal Production* 75:111-113.
- Kerslake JI, Hely FS, Maclennan G, Lindsay S, Behrent MR, Campbell AW 2012. Does castration status have an effect on the quality of lamb meat? *Proceedings of the New Zealand Society of Animal Production* 72: 144–146.
- Schreurs NM 2013. BRIEF COMMUNICATION: Comparison of castrate and entire ram-lambs for meat quality and skatole in the fat. *Proceedings of the New Zealand Society of Animal Production* 73: 68-70.
- SAS Institute Inc. 2011. SAS/STAT® 9.3 User's Guide. Cary, NC: SAS Institute Inc.